

MATHEMATICAL TRIPOS Part III

Thursday 27 May, 2004 1.30 to 4.30

PAPER 2

TOPICS IN GROUP THEORY

*Attempt **THREE** questions.*

*There are **six** questions in total.*

The questions carry equal weight.

**You may not start to read the questions
printed on the subsequent pages until
instructed to do so by the Invigilator.**

1 Let G be a finite group.

(a) Define the Frattini subgroup $\Phi(G)$ and Fitting subgroup $F(G)$ of G . Prove that $F(G)$ exists.

(b) Show that $\Phi(G) \leq F(G)$.

(c) If G is soluble prove that $F(G)$ contains its own centraliser.

(d) Let G be a finite group. The *socle* of G is the subgroup $\text{Soc}(G)$ generated by all the minimal normal subgroups of G . Show that $F(G) \leq C_G(\text{Soc}(G))$.

(e) Prove that $F(G/\Phi(G)) = F(G)/\Phi(G)$.

2 Write an essay on soluble groups. You should include a treatment of Hall π -subgroups.

3 Let G be a doubly transitive permutation group on a finite set Ω of size n and let N be a minimal normal subgroup of G .

(a) If N is regular show that N is elementary abelian of order $n = p^d$ for some prime number p and G is a subgroup of $AGL(d, p)$.

(b) If N is imprimitive show that N is regular.

(c) If N is primitive and non-regular show that N is a non-abelian simple group and $N \leq G \leq \text{Aut}(N)$.

(d) If G is 4-transitive on Ω and N is regular, show that $n = 4$ and $G \cong S_4$.

4 (a) Let G be a group acting primitively on a set Ω and suppose that G is generated by $\{A^g : g \in G\}$, where A is an abelian normal subgroup of G_α for some $\alpha \in \Omega$. If N is a normal subgroup of G , show that either $G' \leq N$ or $N \leq G_{[\Omega]}$ (where $G_{[\Omega]}$ is the pointwise stabiliser of Ω).

(b) Prove that the special linear group $SL(n, \mathbb{F})$ of dimension $n \geq 2$ over a field \mathbb{F} , is generated by transvections.

(c) Prove that $PSL(n, \mathbb{F})$ is simple if $n \geq 3$ or if $n = 2$ and $|\mathbb{F}| > 3$.

(d) Outline the modifications necessary to prove the corresponding result for symplectic groups.

5 (a) Describe how to associate a partition-valued function on the set of irreducible polynomials over \mathbb{F}_q to each conjugacy class of $GL(n, q)$ and give an expression for the size of the corresponding centraliser.

(b) Show that the number of conjugacy classes in $GL(n, q)$ is the coefficient of t^n in

$$\prod_{i=1}^{\infty} \frac{1-t^i}{1-qt^i}.$$

(c) Show that the number of unipotent elements in $GL(n, q)$ is q^{n^2-n} .

[You may assume that $\prod_{i=1}^{\infty} \frac{1}{1-st^i} = \sum_{k=0}^{\infty} \frac{s^k t^k}{\phi_k(t)}$ where $\phi_k(t) = \prod_{i=1}^k (1-t^i)$.]

(d) Let $f(t)$ be a monic irreducible quadratic polynomial over \mathbb{F}_q . Show that the number of elements of $GL(2m, q)$ whose minimal polynomial is a power of $f(t)$ is

$$q^{2m^2-2m} |GL(2m, q)| / |GL(m, q^2)|.$$

6 (a) Define the symplectic group $Sp(2m, q)$ and show that

$$|Sp(2m, q)| = q^{m^2} \prod_{i=1}^m (q^{2i} - 1).$$

(b) Let $\Omega = \{1, 2, \dots, n\}$ and let V be the set of all subsets of Ω . With addition defined by symmetric difference,

$$A + B = (A \cup B) \setminus (A \cap B),$$

V can be regarded as a vector space over the field \mathbb{F}_2 of order 2. Show that the map $(,) : V \times V \rightarrow \mathbb{F}_2$ defined by

$$(A, B) = |A \cap B| \pmod{2},$$

is a symmetric bilinear form on V , preserved by the natural action of the symmetric group S_n .

If n is even, show that $(,)$ induces a non-singular alternating bilinear form on $\langle \Omega \rangle^{\perp} / \langle \Omega \rangle$.

(c) Prove that $Sp(4, 2) \cong S_6$.